

University of Groningen

Sorting and function of peroxisomal membrane proteins

Baerends, RJS; Faber, KN; Kiel, JAKW; van der Klei, IJ; Harder, W; Veenhuis, M

Published in:
FEMS Microbiology Reviews

DOI:
[10.1111/j.1574-6976.2000.tb00543.x](https://doi.org/10.1111/j.1574-6976.2000.tb00543.x)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2000

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Baerends, RJS., Faber, KN., Kiel, JAKW., van der Klei, IJ., Harder, W., & Veenhuis, M. (2000). Sorting and function of peroxisomal membrane proteins. *FEMS Microbiology Reviews*, 24(3), 291-301.
<https://doi.org/10.1111/j.1574-6976.2000.tb00543.x>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Sorting and function of peroxisomal membrane proteins

Richard J.S. Baerends, Klaas Nico Faber, Jan A.K.W. Kiel, Ida J. van der Klei,
Wim Harder, Marten Veenhuis *

Eukaryotic Microbiology, Groningen Biomolecular Sciences and Biotechnology Institute, Postbus 14, 9750 AA Haren, The Netherlands

Received 1 June 1999; received in revised form 10 November 1999; accepted 15 December 1999

Abstract

Peroxisomes are subcellular organelles and are present in virtually all eukaryotic cells. Characteristic features of these organelles are their inducibility and their functional versatility. Their importance in the intermediary metabolism of cells is exemplified by the discovery of several inborn, fatal peroxisomal errors in man, the so-called peroxisomal disorders. Recent findings in research on peroxisome biogenesis and function have demonstrated that peroxisomal matrix proteins and peroxisomal membrane proteins (PMPs) follow separate pathways to reach their target organelle. This paper addresses the principles of PMP sorting and summarizes the current knowledge of the role of these proteins in organelle biogenesis and function. © 2000 Federation of European Microbiological Societies. Published by Elsevier Science B.V. All rights reserved.

Keywords: Peroxisome; Membrane protein; Biogenesis; Protein function; Protein sorting

Contents

1. Introduction	291
2. Peroxisome biogenesis	293
2.1. PMPs involved in organelle biosynthesis and proliferation	293
2.2. Sorting of PMPs	294
2.3. Post-translational modifications	295
2.4. Insertion of PMPs	295
3. Peroxisome function	295
3.1. Metabolite and solute transport	296
3.2. ABC transporters	296
4. Concluding remarks	297
Acknowledgements	297
References	297
References (indicated in Table 1)	300

1. Introduction

Microbodies (peroxisomes, glyoxysomes) represent the last class of organelles that have been discovered [1,2]. They are present in virtually all eukaryotic cells and are

characterized by an unprecedented functional versatility, which varies with the organism in which they occur and with the environmental conditions to which cells are exposed. The observation that in man a group of genetically distinct, mostly fatal, inborn peroxisomal disorders exists [3] has strongly stimulated the interest in the biogenesis and function of these organelles. In particular, the use of yeast and Chinese hamster ovary cells as model systems has been instrumental in attempts to characterize the mo-

* Corresponding author. Tel.: +31 (50) 3632176;
Fax: +31 (50) 3632154; E-mail: m.veenhuis@biol.rug.nl

lecular details of peroxisome assembly. Many mutant strains deficient in peroxisome biogenesis (designated *pex* mutants) have been isolated, the corresponding genes (*PEX* genes) cloned and their protein products (termed peroxins) analyzed (unified nomenclature after [4]). These and other studies have revealed that the principles of microbody biogenesis are highly conserved between yeasts and higher eukaryotes, including man. At present, 22 peroxins have been described; most of them are components of the peroxisomal membrane (see Table 1 and Fig. 1).

Recently, in silico screening of the *Saccharomyces cerevisiae* genome database for protein patterns (i.e. containing the peroxisomal targeting signals; PTS1 or PTS2) or gene expression (i.e. induced in response to oleate) is applicable for the identification of new peroxins [5]. In addition, a comparison of transcript profiles (serial analysis of gene expression) from yeast grown under peroxisome-requiring and non-requiring conditions was also performed [6]. However, these studies indicated that the transcripts encoding known peroxins have a low abundance and are hardly or not induced under peroxisome-requiring conditions, and thus have not yet been the expected source to identify new proteins involved in peroxisome biogenesis [6].

Conceptually, the development of peroxisomes involves a series of events that have to be tuned carefully. These include the recruitment of (phospho)lipids, the synthesis, sorting and assembly/activation of matrix and membrane proteins, as well as organelle inheritance and fission. The peroxisomal membrane mainly consists of phosphatidylcholine and phosphatidylethanolamine and largely resembles that of the endoplasmic reticulum (ER) (reviewed in




Fig. 1. Immunocytochemical localization of the PMP Pex2p in methanol-grown *P. pastoris* wild-type cells, using antibodies against Pex2p (in collaboration with Dr. J.M. Cregg). [aldehyde-unicryl-uranyl acetate] Abbreviations: N, nucleus; P, peroxisome; M, mitochondrion. The bar represents 0.5 μ m.

[7]). How the lipid components of these membranes are recruited is still largely unknown. Only recently, the first experimental data became available which suggest that the ER plays a role in this process and requires vesicle trafficking and fusion events [8–12].

The sorting of microbody matrix proteins is now rather well understood. These proteins are encoded by nuclear genes and, after their synthesis in the cytosol, are recognized by specific soluble receptor molecules. The recognition is mediated by specific targeting signals that reside

Table 1
List of membrane-bound peroxins

Peroxin	Molecular mass (kDa)	Characteristics ^a	References
Pex1p	117–143	AAA-type ATPase; contains two AAA domains; interacts with Pex6p	[1–8]
Pex2p	31–52	IPMP; C-terminal Zn-finger motif; N-glycosylated; sorted via ER ^b	[9–17]
Pex3p	42–52	IPMP; <i>pex3Δ</i> cells do not contain ghosts; interacts with Pex19p	[18–21]
Pex4p	21–24	PMP; ubiquitin-conjugating enzyme	[22–24]
Pex6p	112–127	AAA-type ATPase; contains two AAA domains; interacts with Pex1p	[6–8,25–30]
Pex8p	71–81	PMP facing matrix; contains PTS1 and PTS2	[31–33]
Pex9p	45	IPMP	[34]
Pex10p	34–48	IPMP; C-terminal Zn-finger motif	[35–38]
Pex11p	24–31	IPMP; in mammals two isoforms (α/β); α -form contains dilysine motif, binds coatomer/ARF; dimerizes in baker's yeast and trypanosomes	[39–48]
Pex12p	40–46	IPMP; C-terminal degenerate Zn-finger motif	[49–51]
Pex13p	40–43	IPMP; cytosolic SH3 domain; interacts with Pex5p, Pex7p and Pex14p	[52–54]
Pex14p	38–41	PMP; facing cytosol; phosphorylated; interacts with Pex5p, Pex7p, Pex13p, Pex17p	[55–59]
Pex15p	44	IPMP; phosphorylated	[60]
Pex16p	39–44	PMP; facing matrix; N-glycosylated; sorted via ER ^b	[61–63]
Pex17p	23	PMP; facing cytosol; interacts with Pex14p	[64]
Pex19p	33–40	PMP; facing cytosol; farnesylated; in baker's yeast mainly cytosolic; interacts with Pex3p and Pex10p	[65–69]
Pex22p	23	IPMP; C-terminus cytosolic; interacts with Pex4p	[70]

^a(I)PMP: (integral) peroxisomal membrane protein.

^bIn the yeast *Y. lipolytica*.




Fig. 2. Schematic representation of interactions between soluble and membrane-bound peroxins, as demonstrated by two hybrid analysis or co-immunoprecipitation experiments.

either at the extreme C-terminus (PTS1, recognized by the cytosolic receptor Pex5p) or in the N-terminus of the matrix protein (PTS2, recognized by the cytosolic receptor Pex7p). Most likely, binding of the cargo molecule induces a conformational change in the receptor molecule, which allows the cargo/receptor complex to be recognized by a specific docking protein residing at or in the peroxisomal membrane. At present, two putative docking proteins have been identified, namely Pex13p and Pex14p (see Fig. 2) (reviewed in [13,14]). After release of the cargo protein, the receptor molecule is thought to be recycled to the cytosol. In the methylotrophic yeast *Hansenula polymorpha*, recycling of the PTS1 receptor is dependent on the function of a peroxisomal ubiquitin conjugation enzyme, Pex4p [15]. Whether or not the receptor functionally enters the peroxisomal matrix before recycling is still a matter of debate.

Peroxisomal membrane proteins (PMPs) do not use the sorting machinery of matrix proteins to reach their target organelle, but follow alternative pathways. This paper summarizes the current knowledge on this topic and discusses the role of membrane proteins in peroxisome biogenesis and function.

2. Peroxisome biogenesis

2.1. PMPs involved in organelle biosynthesis and proliferation

Upon induction of peroxisome biogenesis, the synthesis of PMPs was shown to precede that of matrix components [16–18]. In contrast to various inducible matrix proteins, genes that encode essential PMPs are expressed during repressing growth conditions (e.g. on glucose) and are only moderately induced, even when peroxisomes are present at high levels (e.g. in methanol-induced *H. polymorpha*) [19]. This characteristic phenomenon suggests that basic levels of these proteins are necessary for organelle maintenance and inheritance. The finding that the levels of various essential membrane proteins do not increase very much relative to the strong increase in membrane

surface could imply that these proteins are present in functional complexes that are donated to newly formed organelles upon peroxisome fission [20]. The major exception to this is Pex11p; in *S. cerevisiae*, the synthesis of Pex11p is strongly induced (over 30-fold) by oleate and, like many peroxisomal matrix enzymes, is controlled by the oleate-inducible transcription factors Pip2p and Oaf1p [6,21,22].

Both in yeast and in other organisms, Pex11p is essential for normal peroxisome proliferation. This function is proposed from the finding that in oleate-induced cells of a *S. cerevisiae* *pex11Δ* strain, only a few enlarged peroxisomes are observed, while in cells overexpressing *PEX11*, enhanced numbers of relatively small organelles are found [23]. Similar phenotypes were reported when *PEX11* was overexpressed in trypanosomes and mammals, implying similar roles for Pex11p in these organisms [23–26]. *S. cerevisiae* and *Trypanosoma brucei* Pex11p were shown to form homodimers [24,27]. Marshall et al. proposed that Pex11p in its 'monomeric' form is active in proliferation, while increasing oxidative metabolism within maturing peroxisomes causes dimer formation, which inhibits further organelle division [27]. In mammals, two Pex11p isoforms have been identified, encoded by *PEX11α* and *PEX11β* [26,28]. Of these, only *PEX11α* was shown to be induced by peroxisome proliferators, such as clofibrate, whereas *PEX11β* was constitutively expressed [26,28,29]. Interestingly, mammalian Pex11p's contain a dilysine motif at their C-termini, which is absent in Pex11p's [26,28]. Cosson and Letourneur have shown that such dilysine motifs may be involved in binding specific coat proteins (coatamers, COP I vesicles) when exposed to the cytosolic face of the membrane and play a role in the generation of vesicles for the retrieval of ER-resident membrane proteins [30]. Indeed, using purified rat peroxisomes, the group of Just showed that the dilysine motif is able to bind the ADP-ribosylation factor (ARF) and coatamer in a strictly guanosine GTP-γS (5'-O-(3-thiotriphosphate))-dependent manner [25]. This result led them to suggest that by ARF- and coatamer-dependent vesiculation of the peroxisomal membrane, peroxisomes divide or generate nucleation centers for new organelles [25]. However, other mechanisms are likely to exist, since the C-

terminal dilysine motif is not conserved in all Pex11p's identified so far. Also, discrepancies exist in the reported locations of different Pex11p's. In *S. cerevisiae*, Pex11p was suggested to be a membrane-associated protein facing the peroxisomal matrix, whereas in rat and trypanosomes, the N- and C-termini of Pex11p are believed to be exposed to the cytosol [24,25,27]. Detailed biochemical studies will be required to solve these apparent contradictions.

Apart from Pex11p, other peroxins may be involved in peroxisome proliferation, namely Pex10p and Pex16p [31,32]. Peroxisome proliferation can also be regulated by varying the *PEX3* expression level in *H. polymorpha* [33,34]. In constructed conditional mutants of *H. polymorpha* in which the Pex3p levels can be modulated, the size/number of peroxisomes ranged from one or a few large organelles to highly enhanced numbers of small organelles relative to lowered or enhanced Pex3p levels [33,34]. This induction of peroxisome proliferation can occur independent from peroxisome-inducing conditions (extracellular stimuli), since Pex3p overproduced in glucose-grown *H. polymorpha* cells also resulted in proliferation of very small peroxisomes [33]. Essentially, similar results were obtained with *H. polymorpha* Pex14p [35].

Pex deletion strains characteristically contain peroxisomal membrane remnants ('ghosts') in which various peroxisomal membrane components accumulate. The major exception is *pex3Δ*, since in the absence of Pex3p, no peroxisomal remnants were detected [36,37]. Recently, a predominantly cytosolic peroxin, Pex19p, was characterized which was shown to interact with Pex3p (Fig. 2) [38–40]. Also in *pex19* deletion strains, membrane remnants were either completely absent (in *S. cerevisiae*) or morphologically distinct from the late remnants found in other *pex* mutants (in *Pichia pastoris*) [38,39], suggesting that these two proteins are involved in an early stage of peroxisome biogenesis.

2.2. Sorting of PMPs

Heterologous expression experiments have demonstrated that the sorting machinery for PMPs is conserved [24,41–45]. Like peroxisomal matrix proteins, PMPs are encoded by nuclear genes and synthesized on free polyosomes [46–48]. Consequently, they may insert directly from the cytosol into the peroxisomal membrane. Evidence for this came from both in vitro (i.e. mammalian PMP22 and Pex2p) [49,50] and in vivo insertion experiments (i.e. PMP70) [51]. However, other sorting mechanisms may also exist. This can be concluded from the recent findings that several peroxins undergo N- or O-linked glycosylation, thus suggesting that these proteins are sorted to peroxisomes via the ER. The most convincing result in this respect is that in the yeast *Yarrowia lipolytica*, peroxisomal Pex2p and Pex16p were shown to be N-glycosylated. In this yeast, the secretory pathway and peroxisome biogenesis seem to be in part converging

[12,52]. At present, in no other organism, neither N-glycosylation (e.g. human Pex16p) [53,54] nor convergence of the secretory pathway and peroxisome biogenesis have been observed making *Y. lipolytica* an ideal organism to study the process of sorting of PMPs.

Other studies also indicate that the ER may play an essential role in peroxisome biogenesis, although none of them is as yet conclusive. Firstly, Elgersma and coworkers demonstrated that radioactive-labeled mannose was bound to overproduced Pex15p, indicating that the protein is O-mannosylated in *S. cerevisiae* [55]. Additionally, deletion of 30 amino acids from the C-terminus of Pex15p resulted in mislocalization of the truncated protein to the ER [55]. Secondly, we demonstrated that the first 16 N-terminal amino acids of *H. polymorpha* Pex3p are able to direct a reporter protein to the ER [36], suggesting that Pex3p, like Pex2p, Pex15p and Pex16p, is sorted to peroxisomes via the ER. Also peroxisome biogenesis in *H. polymorpha* cells was partially inhibited by brefeldin A, a fungal toxin that interferes with coated vesicle formation. Newly synthesized PMPs (i.e. Pex3p) as well as certain peroxisomal matrix proteins reversibly accumulated in or at the ER [9].

Targeting signals of PMPs (termed mPTSs) have first been identified in *Candida boidinii* PMP47. PMP47 is an integral component of the peroxisomal membrane and a member of the family of mitochondrial solute carriers, containing six transmembrane domains. The mPTS is present at an internal position in a loop between membrane-spanning domains 4 and 5 and contains a stretch of positively charged amino acids [56]. In addition, the mPTS of rat PMP70 was delineated to an internal region around amino acids 190–235 [51]. Subsequently, comparable mPTSs have been identified in Pex3p and Pex15p [37,55,57].

Some general characteristics of the described mPTSs can be defined. In all cases, the mPTS is in close proximity of (a) hydrophobic region(s). The mPTS sequences are clearly distinct from the two targeting sequences so far characterized for peroxisomal matrix proteins, the C-terminal PTS1 and the N-terminal PTS2 (reviewed in [13]). Based on the proposed topologies of PMP47, Pex3p and Pex15p in the peroxisomal membrane, their mPTSs are exposed to the matrix side of the peroxisomal membrane. Furthermore, its precise location is fairly arbitrary, since they have been found in the N-terminal (Pex3p), C-terminal (Pex15p) and an internal (PMP47) position within the primary sequence of the protein.

It may be concluded that distinct classes of mPTSs exist for proteins of different functions. One of these (i.e. in PMP22, PMP47 and PMP70) directs functional proteins (e.g. solute carriers) from the cytosol directly to the peroxisomal membrane. Alternatively, proteins involved in organelle biosynthesis (i.e. Pex2p, Pex3p, Pex15p and Pex16p) may contain sequences directing them to the peroxisome via the ER (Fig. 3).




Fig. 3. Hypothetical pathways of the PMP sorting. Membrane proteins are synthesized on free polysomes in the cytosol. The model discriminates between membrane proteins essential for the biogenesis of organelles (PMBs) and those involved in the organellar function (PMFs, e.g. transporters). PMBs (e.g. Pex2p, Pex3p, Pex15p and Pex16p) may initially be transported to the ER and, eventually vesicle-mediated, be sorted to the growing peroxisome; PMFs may be transported directly to the organelle. Adapted from [9].

At present, no components have been identified which are involved in the recognition of the mPTSs and the subsequent insertion of PMPs into the membrane.

2.3. Post-translational modifications

Several PMPs undergo post-translational modifications. Of these, *H. polymorpha* Pex14p and *S. cerevisiae* Pex15p have been shown to be phosphorylated. However, the functional relevance of these modifications is not yet understood [55,58]. As discussed before, also glycosylation has been observed (see Section 2.2). Human and *S. cerevisiae* Pex19p are farnesylated, a modification that may promote interaction of the protein with the peroxisomal membrane and, furthermore, has been shown to be essential for the proper functioning of the protein in peroxisome biogenesis [38,59]. Apart from membrane anchoring, farnesylation can facilitate protein–protein interactions [60]. This might in fact represent the most probable function in Pex19p since deletion of the farnesylation site strongly affects the interaction with the PMP Pex3p [38]. However, since farnesylation of *P. pastoris* Pex19p is questionable [39], the importance of this modification for peroxisome biogenesis in general needs further investigation. On the other hand, in the C-termini of yeast Pex19p's, the tetrapeptide cysteine–lysine–glutamine–glutamine (CKQQ) is conserved; this sequence was shown to be the best substrate for S-farnesyl transferase, the first enzyme involved in farnesylation [61].

2.4. Insertion of PMPs

In a series of in vitro experiments using purified mammalian peroxisomes, it has been shown that the insertion mechanism of the PMPs Pex2p, PMP22 and PMP70 proceeds in two distinct steps; namely (1) binding to the peroxisomal membrane and (2) integration of the protein into the membrane [49–51]. The insertion of PMP22 and PMP70 did not require ATP or GTP, whereas insertion of Pex2p did require ATP-binding, but not hydrolysis [49–51]. Also, insertion of PMP22 and PMP70 is dependent on (a) proteinaceous component(s) at the peroxisomal membrane which is insensitive to *N*-ethylmaleimide. Cytosolic factors were shown to be involved in the binding and/or insertion of PMP22 and PMP70 [51,62]. These factors most probably represent molecular chaperones or polypeptide chain-binding proteins stabilizing the hydrophobic regions of the proteins thus allowing them to properly insert into the peroxisomal membrane [62]. On the other hand, these factors could represent cytosolic receptors involved in the targeting of the membrane proteins to their target organelle.

3. Peroxisome function

As already predicted from the, often inducible, functional versatility, the peroxisomal membrane must be able to readily adapt in order to allow new metabolic processes to

proceed. A number of the functions that are attributed to the peroxisomal membrane are listed below.

1. The peroxisomal membrane contains several enzymes involved in lipid metabolism (reviewed in [7]).
2. Systems must exist for the recruitment of (phospho)lipids for membrane assembly [63].
3. Association of peroxisomes with the cytoskeleton was shown to be involved in peroxisome morphology, movement and possibly segregation [64–66].
4. The membranes harbor mechanisms that specify recognition and insertion of membrane proteins or import of peroxisomal matrix proteins (e.g. Pex13p, Pex14p, Pex18p and Pex21p) [13,67].
5. Selective degradation of peroxisomes is specified by membrane component(s) [68].
6. Various transport mechanisms (e.g. carriers, transporters) must exist to allow passage of substrates, metabolic intermediates and products across the peroxisomal membrane (reviewed in [69]) as well as cofactors for the peroxisomal enzymes (e.g. FAD, NADH).

3.1. Metabolite and solute transport

The peroxisomal membrane is impermeable in vivo. This was evident from the observation that peroxisomes of *H. polymorpha* are acidic, which suggests that a proton gradient must exist across the peroxisomal membrane [70,71]. As a consequence of this impermeability, various small molecules like NADH, NADPH and acetyl-CoA have to be transported actively. To this purpose, several metabolite shuttles across the membrane have been proposed (e.g. the glycerol-3-phosphate/dihydroxyacetone phosphate and the malate/aspartate shuttles). Instead of transporting the compound itself, they mediate the transfer of metabolic intermediates, which are converted back to the original compound after transport across the membrane [69,72,73]. However, the essential carriers/transporters in these shuttles which transport the intermediates

across the peroxisomal membrane have not yet been identified [20,69].

Among the PMPs identified so far, several members are thought to carry out solute transport (see Table 2). The *C. boidinii* PMP47 gene encodes a 47-kDa integral PMP which is homologous to mitochondrial solute carriers present in the mitochondrial inner membrane (e.g. ADP/ATP exchanger) [41,74]. The N- and C-termini of PMP47 were shown to be exposed to the peroxisomal matrix [75]. Thus, the proposed topology is inverted in comparison to mitochondrial carriers, because their termini extend into the mitochondrial intermembrane space [76]. Deletion of PMP47 in *C. boidinii* wild-type cells causes a defect in the transport and folding of the peroxisomal enzyme dihydroxyacetone synthase (DHAS), whereas other peroxisomal matrix proteins were normally sorted and active. Since deletion of PMP47 in peroxisome-deficient strains did not result in reduced activity of DHAS, it is likely that PMP47 does not catalyze folding directly but is involved in the transport of the DHAS cofactor, thiamine pyrophosphate [77]. Recently, putative human and mouse homologues of *C. boidinii* PMP47, designated PMP34, have been identified. Similar to PMP47, six transmembrane domains were predicted in these proteins [78]. However, the precise function of PMP34 is not yet known.

In addition to the PMP34/PMP47 proteins, another peroxisomal member of the mitochondrial solute carrier family has recently been identified in rabbit, namely a 53-kDa peroxisomal Ca^{2+} -dependent solute carrier [79]. The N-terminus of this protein is exposed to the cytosol and is able to bind Ca^{2+} . Based on the prediction of the presence of six transmembrane domains in the protein and similarity to mitochondrial solute carriers, the authors proposed that the N- and C-termini face the cytosol. Again, the function of the protein is unknown [79].

3.2. ABC transporters

Four ATP-binding cassette (ABC) transporters have

Table 2
PMPs (other than peroxins)

PMP	Molecular mass (kDa)	Characteristics	Reference ^a
PMP20	20	C-terminal PTS1; induced on methanol in yeast	[100–102]
PMP22	22	Pore-forming structure	[103]
PMP34	34	Homologous to mitochondrial carrier family; PMP47 homolog	[78]
PMP47	47	Homologous to mitochondrial carrier family; involved in transport/folding of DHAS	[41,77]
CDSC	53	Ca^{2+} -dependent solute carrier	[79]
PMP70	75	ABC transporter; C-terminal ATP-binding domain; heterodimer	[82,83]
PMP70R	73	ABC transporter; C-terminal ATP-binding domain; heterodimer	[85–87]
ALDP	75	ABC transporter; C-terminal ATP-binding domain; heterodimer	[80,81]
ALDRP	80	ABC transporter; C-terminal ATP-binding domain; heterodimer	[84]
Pxa1p	100	ABC transporter; C-terminal ATP-binding domain; heterodimer (Pxa2p)	[91–94]
Pxa2p	97	ABC transporter; C-terminal ATP-binding domain; heterodimer (Pxa1p)	[91–94]

^aSee references.

been identified in the peroxisomal membrane in mammals: the adrenoleukodystrophy (ALD) protein (ALDP) [80,81], a 70-kDa PMP (PMP70) [82,83], an ALD-related protein (ALDRP) [84] and the PMP70-related protein (PMP70R) [85–87]. ABC transporters are members of the superfamily of membrane proteins involved in transport of various compounds across membranes, ranging from ions to peptides [88]. The transporters are found either as complete transporters (generally 12 transmembrane domains) or as half transporters (generally six transmembrane domains) which can homo- or heterodimerize to form an active transporter [88]. The peroxisomal ABC transporters are all half transporters, which are composed of two functional domains: a transmembrane domain with multiple (six) transmembrane segments and a nucleotide-binding domain with Walker A and B consensus motifs [89] in the C-terminus.

Mutations in the ALD gene cause X-linked ALD, a disorder in peroxisomal β -oxidation resulting in the accumulation of very long chain fatty acids [80]. Although mutations in PMP70 genes of Zellweger syndrome patients have been identified, the specific role of mutant PMP70 in this disease is still speculative [83,90]. ALDRP is most similar to ALDP, whereas PMP70R is most similar to PMP70 [84,85]. The function of both proteins is unknown.

In the genome of the yeast *S. cerevisiae*, only two peroxisomal ABC half transporters have been identified, namely Pxa1p and Pxa2p. These proteins are highly similar to PMP70 and ALDP and were shown to form a heterodimer [91–94]. *S. cerevisiae* Pxa1p and Pxa2p are involved in the transport of long chain fatty acids into peroxisomes [91–93].

Based on the observations in baker's yeast, the mammalian peroxisomal ABC transporters were suggested to mediate transport of fatty acids or enzymes required for β -oxidation into the peroxisomal matrix. The formation of homodimers or different heterodimers by various combinations of the four half transporters could provide different transport facilities towards fatty acids of different saturation states and/or lengths into peroxisomes [91–93]. However, the specific transport function(s) of and physical interaction(s) between the various mammalian or yeast peroxisomal ABC transporters remain to be elucidated. Also, for neither of the half transporters has the topology in the peroxisomal membrane been established [95].

4. Concluding remarks

In the past few years, our knowledge on the biogenesis of peroxisomes has significantly advanced. In particular much has been learned about the mechanisms of matrix protein import which involve novel proteins and new principles. However, major gaps in our knowledge still exist, for instance (1) do PTS receptors functionally enter the peroxisomal matrix and how are they recycled, (2) what

is the actual translocation machinery, (3) are protein import and membrane development coupled processes and (4) is the ER involved in sorting of a specific subset of PMPs and/or phospholipids?

The progress of this work would strongly benefit from the development of an in vitro import system. Although some progress has been made in this field using an in vitro membrane protein insertion assay, a reliable matrix protein import system, in particular for yeast, is not yet available. These studies could also shed light on the basic problem why some proteins can be imported in folded and oligomeric forms (e.g. thiolase) [96] where others are not (e.g. alcohol oxidase) [97].

Another fundamental problem that requires a thorough analysis is the suggestion that peroxisomes may form de novo [5,54,98]. This hypothesis is mainly based on the kinetics of the re-appearance of peroxisomes in specific mutant cells which lack peroxisomal ghosts but in which the originally defective protein is reactivated/reintroduced. As, strictly spoken, de novo synthesis conflicts with the dogma that 'membranes arise from membranes', the alternative explanation is that re-assembly of peroxisomes in transformed *pex* mutants that originally lacked ghosts initiates at the ER. It can be envisaged that upon reintroduction of the corresponding protein, e.g. Pex3p, a peroxisomal import site can be formed at the docking site of Pex3p at the ER, that can act as a pre-peroxisome with the potential to develop into a normal organelle. In line with this reasoning, the ER, or specific regions of the ER, should contain specific docking sites for such peroxins.

Finally, one major line of research that urgently requires exploration concerns peroxisome function. The genetic studies of the group of Tabak have provided evidence that the peroxisomal membrane is not permeable for specific metabolites in vivo [69]. Also, the function of peroxisomes in the metabolism of various substrates (e.g. fatty acids, methanol, primary amines and D-amino acids) predicts that specific transport systems must exist. One of the challenges of the near future is to isolate these carriers and test their function in reconstituted membrane vesicles. Attempts using yeast peroxisomes for in vitro studies failed so far, since the purified organelles are invariably leaky after isolation. Since the protein causing this leakage phenomenon has been identified [99], this aspect can also be addressed.

Acknowledgements

We thank Ineke Keizer-Gunnink for preparing Fig. 1, Florian Salomons for preparing Figs. 2 and 3.

References

- [1] Rhodin, J. (1954) Correlation of ultrastructural organization and

- function in normal and experimentally changed proximal convoluted cells of the mouse kidney. Ph.D. thesis, Stockholm.
- [2] De Duve, C. and Baudhuin, P. (1966) Peroxisomes (microbodies and related particles). *Physiol. Rev.* 46, 323–357.
 - [3] van den Bosch, H., Schutgens, R.B.H., Wanders, R.J.A. and Tager, J.M. (1992) Biochemistry of peroxisomes. *Annu. Rev. Biochem.* 61, 157–197.
 - [4] Distel, B., Erdmann, R., Gould, S.J., Blobel, G., Crane, D.I., Cregg, J.M., Dodt, G., Fujiki, Y., Goodman, J.M., Just, W.W., Kiel, J.A.K.W., Kunau, W.-H., Lazarow, P.B., Mannaerts, G.P., Moser, H.W., Osumi, T., Rachubinski, R.A., Roscher, A., Subramani, S., Tabak, H.F., Tsukamoto, T., Valle, D., van der Klei, I., van Veldhoven, P.P. and Veenhuis, M. (1996) A unified nomenclature for peroxisome biogenesis factors. *J. Cell Biol.* 135, 1–3.
 - [5] Geraghty, M.T., Bassett, D., Morrell, J.C., Gatto Jr., G.J., Bai, J., Geisbrecht, B.V., Hieter, P. and Gould, S.J. (1999) Detecting patterns of protein distribution and gene expression in silico. *Proc. Natl. Acad. Sci. USA* 96, 2937–2942.
 - [6] Kal, A.J., van Zonneveld, A.J., Benes, V., van den Berg, M., Groot Koerkamp, M., Albermann, K., Strack, N., Ruijter, J.M., Richter, A., Dujon, B., Ansoorge, W. and Tabak, H.F. (1999) Dynamics of gene expression revealed by comparison of serial analysis of gene expression transcript profiles from yeast grown on two different carbon sources. *Mol. Biol. Cell* 10, 1859–1872.
 - [7] van Veldhoven, P.P. and Mannaerts, G.P. (1994) Assembly of the peroxisomal membrane. In: *Subcellular Biochemistry, Membrane Biogenesis*, Vol. 22 (Maddy, A.H. and Harris, J.R., Eds.), pp. 231–261. Plenum Press, New York.
 - [8] Baerends, R.J.S., Rasmussen, S.W., Hilbrands, R.E., van der Heide, M., Faber, K.N., Reuvekamp, P.T.W., Kiel, J.A.K.W., Cregg, J.M., van der Klei, I.J. and Veenhuis, M. (1996) The *Hansenula polymorpha* PER9 gene encodes a peroxisomal membrane protein essential for peroxisome assembly and integrity. *J. Biol. Chem.* 271, 8887–8894.
 - [9] Salomons, F.A., van der Klei, I.J., Kram, A.M., Harder, W. and Veenhuis, M. (1997) Brefeldin A interferes with peroxisomal protein sorting in the yeast *Hansenula polymorpha*. *FEBS Lett.* 411, 133–139.
 - [10] Elgersma, Y., Kwast, L., van den Berg, M., Snyder, W.B., Distel, B., Subramani, S. and Tabak, H.F. (1997) Overexpression of Pex15p, a phosphorylated peroxisomal integral membrane protein required for peroxisome assembly in *S. cerevisiae*, causes proliferation of the endoplasmic reticulum membrane. *EMBO J.* 16, 7326–7341.
 - [11] Faber, K.N., Heyman, J.A. and Subramani, S. (1998) Two AAA family peroxins, PpPex1p and PpPex6p, interact with each other in an ATP-dependent manner and are associated with different subcellular membranous structures distinct from peroxisomes. *Mol. Cell. Biol.* 18, 936–943.
 - [12] Titorenko, V.I. and Rachubinski, R.A. (1998) Mutants of the yeast *Yarrowia lipolytica* defective in protein exit from the endoplasmic reticulum are also defective in peroxisome biogenesis. *Mol. Cell. Biol.* 18, 2789–2803.
 - [13] Erdmann, R., Veenhuis, M. and Kunau, W.-H. (1997) Peroxisomes: organelles at the crossroads. *Trends Cell Biol.* 7, 400–407.
 - [14] Subramani, S. (1998) Components involved in peroxisome import, biogenesis, proliferation, turnover and movement. *Physiol. Rev.* 78, 171–188.
 - [15] van der Klei, I.J., Hilbrands, R.E., Kiel, J.A.K.W., Rasmussen, S.W., Cregg, J.M. and Veenhuis, M. (1998) The ubiquitin-conjugating enzyme Pex4p of *Hansenula polymorpha* is required for efficient functioning of the PTS1 import machinery. *EMBO J.* 17, 3608–3618.
 - [16] Lüers, G., Beier, K., Hashimoto, T., Fahimi, H.D. and Volkl, A. (1990) Biogenesis of peroxisomes: sequential biosynthesis of the membrane and matrix proteins in the course of hepatic regeneration. *Eur. J. Cell Biol.* 52, 175–184.
 - [17] Veenhuis, M. and Goodman, J.M. (1990) Peroxisomal assembly: membrane proliferation precedes the induction of the abundant matrix proteins in the methylotrophic yeast *Candida boidinii*. *J. Cell Sci.* 96, 583–590.
 - [18] Erdmann, R. and Blobel, G. (1995) Giant peroxisomes in oleic acid-induced *Saccharomyces cerevisiae* lacking the peroxisomal membrane protein Pmp27p. *J. Cell Biol.* 128, 509–523.
 - [19] Sulter, G.J., Looyenga, L., Veenhuis, M. and Harder, W. (1990) Occurrence of peroxisomal membrane proteins in methylotrophic yeasts grown under different conditions. *Yeast* 6, 35–43.
 - [20] van der Klei, I.J. and Veenhuis, M. (1997) Yeast peroxisomes: function and biogenesis of a versatile cell organelle. *Trends Microbiol.* 5, 502–509.
 - [21] Rottensteiner, H., Kal, A.J., Filipits, M., Binder, M., Hamilton, B., Tabak, H.F. and Ruis, H. (1996) Pip2p: a transcriptional regulator of peroxisome proliferation in the yeast *Saccharomyces cerevisiae*. *EMBO J.* 15, 2924–2934.
 - [22] Karpichev, I.V. and Small, G.M. (1998) Global regulatory functions of Oaf1p and Pip2p (Oaf2p), transcription factors that regulate genes encoding peroxisomal proteins in *Saccharomyces cerevisiae*. *Mol. Cell Biol.* 18, 6560–6570.
 - [23] Marshall, P.A., Krimkevich, Y.I., Lark, R.H., Dyer, J.M., Veenhuis, M. and Goodman, J.M. (1995) Pmp27 promotes peroxisomal proliferation. *J. Cell Biol.* 129, 345–355.
 - [24] Lorenz, P., Maier, A.G., Baumgart, E., Erdmann, R. and Clayton, C. (1998) Elongation and clustering of glycosomes in *Trypanosoma brucei* overexpressing the glycosomal Pex11p. *EMBO J.* 17, 3542–3555.
 - [25] Passreiter, M., Anton, M., Lay, D., Frank, R., Harter, C., Wieland, F.T., Gorgas, K. and Just, W.W. (1998) Peroxisome biogenesis: involvement of ARF and coatamer. *J. Cell Biol.* 141, 373–383.
 - [26] Schrader, M., Reuber, B.E., Morrell, J.C., Jimenez-Sanchez, G., Obie, C., Stroh, T.A., Valle, D., Schroer, T.A. and Gould, S.J. (1998) Expression of *PEX11 β* mediates peroxisome proliferation in the absence of extracellular stimuli. *J. Biol. Chem.* 273, 29607–29614.
 - [27] Marshall, P.A., Dyer, J.M., Quick, M.E. and Goodman, J.M. (1996) Redox-sensitive homodimerization of Pex11p: a proposed mechanism to regulate peroxisomal division. *J. Cell Biol.* 135, 123–137.
 - [28] Abe, I., Okumoto, K., Tamura, S. and Fujiki, Y. (1998) Clofibrate-inducible, 28-kDa peroxisomal integral membrane protein is encoded by *PEX11*. *FEBS Lett.* 431, 468–472.
 - [29] Abe, I. and Fujiki, Y. (1998) cDNA cloning and characterization of a constitutively expressed isoform of the human peroxin Pex11p. *Biochem. Biophys. Res. Commun.* 252, 529–533.
 - [30] Cosson, P. and Letourneur, F. (1994) Coatamer interaction with di-lysine endoplasmic reticulum motifs. *Science* 263, 1629–1631.
 - [31] Tan, X., Waterham, H.R., Veenhuis, M. and Cregg, J.M. (1995) The *Hansenula polymorpha* PER8 gene encodes a novel peroxisomal integral membrane protein involved in proliferation. *J. Cell Biol.* 128, 307–319.
 - [32] Eitzen, G.A., Szilard, R.K. and Rachubinski, R.A. (1997) Enlarged peroxisomes are present in oleic acid-grown *Yarrowia lipolytica* overexpressing the *PEX16* gene encoding an intraperoxisomal peripheral membrane peroxin. *J. Cell Biol.* 137, 1265–1278.
 - [33] Baerends, R.J.S., Salomons, F.A., Faber, K.N., Kiel, J.A.K.W., van der Klei, I.J. and Veenhuis, M. (1997) Deviant Pex3p levels affect normal peroxisome formation in *Hansenula polymorpha*: high steady-state levels of the protein fully abolish matrix protein import. *Yeast* 13, 1437–1448.
 - [34] Baerends, R.J.S., Salomons, F.A., Kiel, J.A.K.W., van der Klei, I.J. and Veenhuis, M. (1998) Deviant Pex3p levels affect normal peroxisome formation in *Hansenula polymorpha*: a sharp increase of the protein level induces the proliferation of numerous, small protein-import competent peroxisomes. *Yeast* 13, 1449–1463.
 - [35] Komori, M., Rasmussen, S.W., Kiel, J.A.K.W., Baerends, R.J.S., Cregg, J.M., van der Klei, I.J. and Veenhuis, M. (1997) The *Hansenula polymorpha* PEX14 gene encodes a novel peroxisomal membrane protein essential for peroxisome biogenesis. *EMBO J.* 16, 44–53.
 - [36] Baerends, R.J.S., Rasmussen, S.W., Hilbrands, R.E., van der Heide, M., Faber, K.N., Reuvekamp, P.T.W., Kiel, J.A.K.W., Cregg, J.M., van der Klei, I.J. and Veenhuis, M. (1996) The *Hansenula polymorpha*

- PER9* gene encodes a peroxisomal membrane protein essential for peroxisome assembly and integrity. *J. Biol. Chem.* 271, 8887–8894.
- [37] Wiemer, E.A.C., Lüers, G.H., Faber, K.N., Wenzel, T., Veenhuis, M. and Subramani, S. (1996) Isolation and characterization of Pas2p, a peroxisomal membrane protein essential for peroxisome biogenesis in the methylotrophic yeast *Pichia pastoris*. *J. Biol. Chem.* 271, 18973–18980.
 - [38] Götte, K., Girzalsky, W., Linkert, M., Baumgart, E., Kammerer, S., Kunau, W.-H. and Erdmann, R. (1998) Pex19p, a farnesylated protein essential for peroxisome biogenesis. *Mol. Cell. Biol.* 18, 616–628.
 - [39] Snyder, W.B., Faber, K.N., Wenzel, T.J., Koller, A., Lüers, G.H., Rangell, L., Keller, G.A. and Subramani, S. (1999) Pex19p interacts with Pex3p and Pex10p and is essential for peroxisome biogenesis in *Pichia pastoris*. *Mol. Biol. Cell* 10, 1745–1761.
 - [40] Soukupova, M., Sprenger, C., Gorgas, K., Kunau, W.-H. and Dodt, G. (1999) Identification and characterization of the human peroxin PEX3. *Eur. J. Cell Biol.* 78, 357–374.
 - [41] McCammon, M.T., Dowds, C.A., Orth, K., Moomaw, C.R., Slaughter, C.A. and Goodman, J.M. (1990) Sorting of peroxisomal membrane protein PMP47 from *Candida boidinii* into peroxisomal membranes of *Saccharomyces cerevisiae*. *J. Biol. Chem.* 265, 20098–20105.
 - [42] Sulter, G.J., Waterham, H.R., Vrieling, E.G., Goodman, J.M., Harder, W. and Veenhuis, M. (1993) Expression and targeting of a 47 kDa integral peroxisomal membrane protein of *Candida boidinii* in wild-type and a peroxisome-deficient mutant of *Hansenula polymorpha*. *FEBS Lett.* 315, 211–216.
 - [43] Kiel, J.A.K.W., Keizer-Gunnink, I.K., Krause, T., Komori, M. and Veenhuis, M. (1995) Heterologous complementation of peroxisome function in yeast: the *Saccharomyces cerevisiae* *PAS3* gene restores peroxisome biogenesis in a *Hansenula polymorpha* *per9* disruption mutant. *FEBS Lett.* 377, 434–438.
 - [44] Sakai, Y., Marshall, P.A., Saiganji, A., Takabe, K., Saiki, H., Kato, N. and Goodman, J.M. (1995) The *Candida boidinii* peroxisomal membrane protein Pmp30 has a role in peroxisomal proliferation and is functionally homologous to Pmp27 from *Saccharomyces cerevisiae*. *J. Bacteriol.* 177, 6773–6781.
 - [45] Elgersma, Y., Kwast, L., Klein, A., Voorn-Brouwer, T., van den Berg, M., Metzger, B., America, T., Tabak, H.F. and Distel, B. (1996) The SH3 domain of the *Saccharomyces cerevisiae* peroxisomal membrane protein Pex13p functions as a docking site for Pex5p, a mobile receptor for the import of PTS1-containing proteins. *J. Cell Biol.* 135, 97–109.
 - [46] Fujiki, Y., Rachubinski, R.A. and Lazarow, P.B. (1984) Synthesis of a major integral membrane polypeptide of rat liver peroxisomes on free polysomes. *Proc. Natl. Acad. Sci. USA* 81, 7127–7131.
 - [47] Suzuki, Y., Orii, T., Takiguchi, M., Mori, M., Hijikata, M. and Hashimoto, T. (1987) Biosynthesis of membrane polypeptides of rat liver peroxisomes. *J. Biochem. (Tokyo)* 101, 491–496.
 - [48] Just, W.W. and Hartl, F.-U. (1987) Peroxisomes in Biology and Medicine (Fahimi, H.D. and Sie, H., Eds.), pp. 402–416. Springer-Verlag, Heidelberg.
 - [49] Diestelkötter, P. and Just, W.W. (1993) In vitro insertion of the 22-kDa peroxisomal membrane protein into isolated rat liver peroxisomes. *J. Cell Biol.* 123, 1717–1725.
 - [50] Just, W.W. and Diestelkötter, P. (1996) Protein insertion into the peroxisomal membrane. *Ann. N.Y. Acad. Sci.* 804, 60–75.
 - [51] Imanaka, T., Shiina, Y., Takano, T., Hashimoto, T. and Osumi, T. (1996) Insertion of the 70-kDa peroxisomal membrane protein into peroxisomal membranes in vivo and in vitro. *J. Biol. Chem.* 271, 3706–3713.
 - [52] Titorenko, V.I., Ogrydziak, D.M. and Rachubinski, R.A. (1997) Four distinct secretory pathways serve protein secretion, cell surface growth, and peroxisome biogenesis in the yeast *Yarrowia lipolytica*. *Mol. Cell. Biol.* 17, 5210–5226.
 - [53] Honsho, M., Tamura, S., Shimozaawa, N., Suzuki, Y., Kondo, N. and Fujiki, Y. (1998) Mutation in *PEX16* is causal in the peroxisome-deficient Zellweger syndrome of complementation group D. *Am. J. Hum. Genet.* 63, 1622–1630.
 - [54] South, S.T. and Gould, S.J. (1999) Peroxisome synthesis in the absence of preexisting peroxisomes. *J. Cell Biol.* 144, 255–266.
 - [55] Elgersma, Y., Kwast, L., van den Berg, M., Snyder, W.B., Distel, B., Subramani, S. and Tabak, H.F. (1997) Overexpression of Pex15p, a phosphorylated peroxisomal integral membrane protein required for peroxisome assembly in *S. cerevisiae*, causes proliferation of the endoplasmic reticulum membrane. *EMBO J.* 16, 7326–7341.
 - [56] Dyer, J.M., McNew, J.A. and Goodman, J.M. (1996) The sorting sequence of the peroxisomal integral membrane protein PMP47 is contained within a short hydrophilic loop. *J. Cell Biol.* 133, 269–280.
 - [57] Kammerer, S., Holzinger, A., Welsch, U. and Roscher, A.A. (1998) Cloning and characterization of the gene encoding the human peroxisomal assembly protein Pex3p. *FEBS Lett.* 429, 53–60.
 - [58] Komori, M., Kiel, J.A.K.W. and Veenhuis, M. (1999) The peroxisomal membrane protein Pex14p of *Hansenula polymorpha* is phosphorylated in vivo. *FEBS Lett.* 457, 397–399.
 - [59] Matsuzono, Y., Kinoshita, N., Tamura, S., Shimozaawa, N., Hama-saki, M., Ghaedi, K., Wanders, R.J.A., Suzuki, Y., Kondo, N. and Fujiki, Y. (1999) Human *PEX19*: cDNA cloning by functional complementation, mutation analysis in a patient with Zellweger syndrome, and potential role in peroxisomal membrane assembly. *Proc. Natl. Acad. Sci. USA* 96, 2116–2121.
 - [60] Zhang, F.L. and Casey, P.J. (1996) Protein prenylation: molecular mechanisms and functional consequences. *Annu. Rev. Biochem.* 65, 241–269.
 - [61] Boutin, J.A., Marande, W., Petit, L., Loynel, A., Desmet, C., Canet, E. and Fauchère, J.-L. (1999) Investigations of S-farnesyl transferase substrate specificity with combinatorial tetrapeptide libraries. *Cell. Signal.* 11, 59–69.
 - [62] Pause, B., Diestelkötter, P., Heid, H. and Just, W.W. (1997) Cytosolic factors mediate protein insertion into the peroxisomal membrane. *FEBS Lett.* 414, 95–98.
 - [63] Wirtz, K.W.A. (1982) Lipid-Protein Interactions (Jost, P. and Griffith, O.H., Eds.), pp. 151–231. Wiley, New York.
 - [64] Schrader, M., Burkhardt, J.K., Baumgart, E., Lüers, G., Spring, H., Völkl, A. and Fahimi, H.D. (1996) Interaction of microtubules with peroxisomes. Tubular and spherical peroxisomes in HepG2 cells and their alterations induced by microtubule-active drugs. *Eur. J. Cell Biol.* 69, 24–35.
 - [65] Huber, C.M., Saffrich, R., Anton, M., Passreiter, M., Ansorge, W., Gorgas, K. and Just, W.W. (1997) A heterotrimeric G protein-phospholipase A₂ signaling cascade is involved in the regulation of peroxisomal motility in CHO cells. *J. Cell Sci.* 110, 2955–2968.
 - [66] Wiemer, E.A.C., Wenzel, T., Deerinck, T.J., Ellisman, M.H. and Subramani, S. (1997) Visualization of the peroxisomal compartment in living mammalian cells: dynamic behavior and association with microtubules. *J. Cell Biol.* 136, 71–80.
 - [67] Purdue, P.E., Yang, X. and Lazarow, P.B. (1998) Pex18p and Pex21p, a novel pair of related peroxins essential for peroxisomal targeting by the PTS2 pathway. *J. Cell Biol.* 143, 1859–1869.
 - [68] Kiel, J.A.K.W., Rechinger, K.B., van der Klei, I.J., Salomons, F.A., Titorenko, V.I. and Veenhuis, M. (1999) The *Hansenula polymorpha* *PDD1* gene product, essential for the selective degradation of peroxisomes, is a homologue of *Saccharomyces cerevisiae* Vps34p. *Yeast* 15, 741–754.
 - [69] Elgersma, Y. and Tabak, H.F. (1996) Proteins involved in peroxisome biogenesis and functioning. *Biochim. Biophys. Acta* 1286, 269–283.
 - [70] Nicolay, K., Veenhuis, M., Douma, A.C. and Harder, W. (1987) A ³¹P NMR study of the internal pH of yeast peroxisomes. *Arch. Microbiol.* 147, 37–41.
 - [71] Waterham, H.R., Keizer-Gunnink, I., Goodman, J.M., Harder, W. and Veenhuis, M. (1990) Immunocytochemical evidence for the acidic nature of peroxisomes of methylotrophic yeasts. *FEBS Lett.* 262, 17–19.

- [72] Baumgart, E., Fahimi, H.D., Stich, A. and Völkl, A. (1996) L-lactate dehydrogenase A4- and A3B isoforms are bona fide peroxisomal enzymes in rat liver. Evidence for involvement in intraperoxisomal NADH reoxidation. *J. Biol. Chem.* 271, 3846–3855.
- [73] van Roermund, C.W., Hetteima, E.H., Kal, A.J., van den Berg, M., Tabak, H.F. and Wanders, R.J.A. (1998) Peroxisomal beta-oxidation of polyunsaturated fatty acids in *Saccharomyces cerevisiae*: isocitrate dehydrogenase provides NADPH for reduction of double bonds at even positions. *EMBO J.* 17, 677–687.
- [74] Jank, B., Habermann, B., Schweyen, R.J. and Link, T.A. (1993) PMP47, a peroxisomal homologue of mitochondrial solute carrier proteins. *Trends Biochem. Sci.* 18, 427–428.
- [75] McCammon, M.T., McNew, J.A., Willy, P.J. and Goodman, J.M. (1994) An internal region of the peroxisomal membrane protein PMP47 is essential for sorting to peroxisomes. *J. Cell Biol.* 124, 915–925.
- [76] Kuan, J. and Saier Jr., M.H.J. (1993) The mitochondrial carrier family of transport proteins: structural, functional, and evolutionary relationships. *Crit. Rev. Biochem. Mol. Biol.* 28, 209–233.
- [77] Sakai, Y., Saiganji, A., Yurimoto, H., Takabe, K., Saiki, H. and Kato, N. (1996) The absence of Pmp47, a putative yeast peroxisomal transporter, causes a defect in transport and folding of a specific matrix enzyme. *J. Cell Biol.* 134, 37–51.
- [78] Wylin, T., Baes, M., Brees, C., Mannaerts, G.P., Fransen, M. and van Veldhoven, P.P. (1998) Identification and characterization of human PMP34, a protein closely related to the peroxisomal integral membrane protein PMP47 of *Candida boidinii*. *Eur. J. Biochem.* 258, 332–338.
- [79] Weber, F.E., Minestrini, G., Dyer, J.H., Werder, M., Boffelli, D., Compassi, S., Wehrli, E., Thomas, R.M., Schulthess, G. and Hauser, H. (1997) Molecular cloning of a peroxisomal Ca^{2+} -dependent member of the mitochondrial carrier superfamily. *Proc. Natl. Acad. Sci. USA* 94, 8509–8514.
- [80] Mosser, J., Douar, A.M., Sarde, C.O., Kioschis, P., Feil, R., Moser, H., Poustka, A.M., Mandel, J.L. and Aubourg, P. (1993) Putative X-linked adrenoleukodystrophy gene shares unexpected homology with ABC transporters. *Nature* 361, 726–730.
- [81] Mosser, J., Lutz, Y., Stoeckel, M.E., Sarde, C.O., Kretz, C., Douar, A.M., Lopez, J., Aubourg, P. and Mandel, J.L. (1994) The gene responsible for adrenoleukodystrophy encodes a peroxisomal membrane protein. *Hum. Mol. Genet.* 3, 265–271.
- [82] Kamijo, K., Taketani, S., Yokota, S., Osumi, T. and Hashimoto, T. (1990) The 70-kDa peroxisomal membrane protein is a member of the Mdr (P-glycoprotein)-related ATP-binding protein superfamily. *J. Biol. Chem.* 265, 4534–4540.
- [83] Gärtner, J., Moser, H. and Valle, D. (1992) Mutations in the 70K peroxisomal membrane protein gene in Zellweger syndrome. *Nat. Genet.* 1, 16–23.
- [84] Lombard-Platet, G., Savary, S., Sarde, C.O., Mandel, J.L. and Chimini, G. (1996) A close relative of the adrenoleukodystrophy (ALD) gene codes for a peroxisomal protein with a specific expression pattern. *Proc. Natl. Acad. Sci. USA* 93, 1265–1269.
- [85] Shani, N., Jimenez-Sanchez, G., Steel, G., Dean, M. and Valle, D. (1997) Identification of a fourth half ABC transporter in the human peroxisomal membrane. *Hum. Mol. Genet.* 6, 1925–1931.
- [86] Holzinger, A., Roscher, A.A., Landgraf, P., Lichtner, P. and Kammerer, S. (1998) Genomic organization and chromosomal localization of the human peroxisomal membrane protein-1-like protein (PXMP1-L) gene encoding a peroxisomal ABC transporter. *FEBS Lett.* 426, 238–242.
- [87] Holzinger, A., Muntau, A., Mayerhofer, P., Kammerer, S., Albet, S., Bugaut, M. and Roscher, A.A. (1998) The mouse gene encoding the peroxisomal membrane protein 1-like protein (PXMP1-L): cDNA cloning, genomic organization and comparative expression studies. *FEBS Lett.* 433, 179–183.
- [88] Higgins, C.F. (1992) ABC transporters: from microorganisms to man. *Annu. Rev. Cell Biol.* 8, 67–113.
- [89] Walker, J.E., Saraste, M., Runswick, M.J. and Gay, N.J. (1982) Distantly related sequences in the α - and β -subunits of ATP synthase, myosin, kinases and other ATP-requiring enzymes and a common nucleotide binding fold. *EMBO J.* 1, 945–981.
- [90] Braverman, N., Dodt, G., Gould, S.J. and Valle, D. (1995) Disorders of peroxisome biogenesis. *Hum. Mol. Genet.* 4, 1791–1798.
- [91] Shani, N., Watkins, P.A. and Valle, D. (1995) *PXAI*, a possible *Saccharomyces cerevisiae* ortholog of the human adrenoleukodystrophy gene. *Proc. Natl. Acad. Sci. USA* 92, 6006–6012.
- [92] Hetteima, E.H., van Roermund, C.W., Distel, B., van den Berg, M., Vilela, C., Rodrigues-Pousada, C., Wanders, R.J.A. and Tabak, H.F. (1996) The ABC transporter proteins Pat1 and Pat2 are required for import of long-chain fatty acids into peroxisomes of *Saccharomyces cerevisiae*. *EMBO J.* 15, 3813–3822.
- [93] Shani, N. and Valle, D. (1996) A *Saccharomyces cerevisiae* homolog of the human adrenoleukodystrophy transporter is a heterodimer of two half ATP-binding cassette transporters. *Proc. Natl. Acad. Sci. USA* 93, 11901–11906.
- [94] Swartzman, E.E., Viswanathan, M.N. and Thorner, J. (1996) The *PAL1* gene product is a peroxisomal ATP-binding cassette transporter in the yeast *Saccharomyces cerevisiae*. *J. Cell Biol.* 132, 549–563.
- [95] Shani, N. and Valle, D. (1998) Peroxisomal ABC transporters. *Methods Enzymol.* 292, 753–776.
- [96] Glover, J.R., Andrews, D.W. and Rachubinski, R.A. (1994) *Saccharomyces cerevisiae* peroxisomal thiolase is imported as a dimer. *Proc. Natl. Acad. Sci. USA* 91, 10541–10545.
- [97] Waterham, H.R., Russell, K.A., Vries, Y. and Cregg, J.M. (1997) Peroxisomal targeting, import, and assembly of alcohol oxidase in *Pichia pastoris*. *J. Cell Biol.* 139, 1419–1431.
- [98] Waterham, H.R., Titorenko, V.I., Swaving, G.J., Harder, W. and Veenhuis, M. (1993) Peroxisomes in the methylotrophic yeast *Hansenula polymorpha* do not necessarily derive from pre-existing organelles. *EMBO J.* 12, 4785–4794.
- [99] Sulter, G.J., Verheyden, K., Mannaerts, G., Harder, W. and Veenhuis, M. (1993) The in vitro permeability of yeast peroxisomal membranes is caused by a 31 kDa integral membrane protein. *Yeast* 9, 733–742.
- [100] Garrard, L.J. and Goodman, J.M. (1989) Two genes encode the major membrane-associated protein of methanol-induced peroxisomes from *Candida boidinii*. *J. Biol. Chem.* 264, 13929–13937.
- [101] Rande-Gil, F., Prieto, J.A. and Sanz, P. (1994) Nucleotide sequence of a putative peroxisomal protein from the yeast *lipomyces kononenkoae*. *FEMS Microbiol. Lett.* 122, 153–158.
- [102] Yamashita, H., Avraham, S., Jiang, S., London, R., van Veldhoven, P.P., Subramani, S., Rogers, R.A. and Avraham, H. (1999) Characterization of human and murine PMP20 peroxisomal proteins that exhibit antioxidant activity in vitro. *J. Biol. Chem.* 274, 29897–29904.
- [103] Kaldi, K., Diestelkötter, P., Stenbeck, G., Auerbach, S., Jäckle, U., Mägert, H.-J., Wieland, F.T. and Just, W.W. (1993) Membrane topology of the 22 kDa integral peroxisomal membrane protein. *FEBS Lett.* 315, 217–222.

References (indicated in Table 1)

- [1] Erdmann, R. et al. (1991) *Cell* 64, 499–510.
- [2] Heyman, J.A. et al. (1994) *J. Cell Biol.* 127, 1259–1273.
- [3] Portsteffen, H. et al. (1997) *Nat. Genet.* 17, 449–452.
- [4] Reuber, B.E. et al. (1997) *Nat. Genet.* 17, 445–448.
- [5] Tamura, S. et al. (1998) *Proc. Natl. Acad. Sci. USA* 95, 4350–4355.
- [6] Tamura, S. et al. (1998) *Biochem. Biophys. Res. Commun.* 245, 883–886.
- [7] Faber, K.N. et al. (1998) *Mol. Cell. Biol.* 18, 936–943.
- [8] Kiel, J.A.K.W. et al. (1999) *Yeast* 15, 1059–1078.

- [9] Tsukamoto, T. et al. (1991) *Nature* 350, 77–81.
- [10] Shimozaawa, N. et al. (1992) *Science* 255, 1132–1134.
- [11] Thieringer, R. and Rietz, C.R.H. (1993) *J. Biol. Chem.* 268, 12631–12636.
- [12] Wilson, G.N. and Bryant, D.D. (1994) *Biochem. Med. Metab. Biol.* 51, 140–148.
- [13] Berteaux-Lecellier, V. et al. (1995) *Cell* 81, 1043–1051.
- [14] Eitzen, G.A. et al. (1996) *J. Biol. Chem.* 271, 20300–20306.
- [15] Liu, X. et al. (1996) *Yeast* 12, 135–143.
- [16] Waterham, H.R. et al. (1996) *Mol. Cell. Biol.* 16, 2527–2536.
- [17] Flaspohler, J.A. et al. (1997) *Mol. Cell. Biol.* 17, 1093–1101.
- [18] Höhfeld, J. et al. (1991) *J. Cell Biol.* 114, 1167–1178.
- [19] Baerends, R.J.S. et al. (1996) *J. Biol. Chem.* 271, 8887–8894.
- [20] Wiemer, E.A.C. et al. (1996) *J. Biol. Chem.* 271, 18973–18980.
- [21] Kammerer, S. et al. (1998) *FEBS Lett.* 429, 53–60.
- [22] Wiebel, F.F. and Kunau, W.-H. (1992) *Science* 359, 73–76.
- [23] Crane, D.I. et al. (1994) *J. Biol. Chem.* 269, 21835–21844.
- [24] van der Klei, I.J. et al. (1998) *EMBO J.* 17, 3608–3618.
- [25] Spong, A.P. and Subramani, S. (1993) *J. Cell Biol.* 123, 535–548.
- [26] Voorn-Brouwer, T. et al. (1993) *Biochim. Biophys. Acta* 1216, 325–328.
- [27] Nuttley, W.M. et al. (1994) *J. Biol. Chem.* 269, 556–566.
- [28] Tsukamoto, T. et al. (1995) *Nat. Genet.* 11, 395–401.
- [29] Yahraus, T. et al. (1996) *EMBO J.* 15, 2914–2923.
- [30] Fukuda, S. et al. (1996) *Am. J. Hum. Genet.* 59, 1210–1220.
- [31] Waterham, H.R. et al. (1994) *J. Cell Biol.* 127, 737–749.
- [32] Liu, H. et al. (1995) *J. Biol. Chem.* 270, 10940–10951.
- [33] Smith, J.J. et al. (1997) *Mol. Cell. Biol.* 17, 2511–2520.
- [34] Eitzen, G.A. et al. (1995) *J. Biol. Chem.* 270, 1429–1436.
- [35] Tan, X. et al. (1995) *J. Cell Biol.* 128, 307–319.
- [36] Kalish, J.E. et al. (1995) *Mol. Cell. Biol.* 15, 6406–6419.
- [37] Okumoto, K. et al. (1998) *Hum. Mol. Genet.* 7, 1399–1405.
- [38] Warren, D.S. et al. (1998) *Am. J. Hum. Genet.* 63, 347–359.
- [39] Moreno, M. et al. (1994) *Yeast* 10, 1447–1457.
- [40] Erdmann, R. and Blobel, G. (1995) *J. Cell Biol.* 128, 509–523.
- [41] Marshall, P.A. et al. (1995) *J. Cell Biol.* 129, 345–355.
- [42] Sakai, Y. et al. (1995) *J. Bacteriol.* 177, 6773–6781.
- [43] Marshall, P.A. et al. (1996) *J. Cell Biol.* 135, 123–137.
- [44] Abe, I. et al. (1998) *FEBS Lett.* 431, 468–472.
- [45] Abe, I. and Fujiki, Y. (1998) *Biochem. Biophys. Res. Comm.* 252, 529–533.
- [46] Lorenz, P. et al. (1998) *EMBO J.* 17, 3542–3555.
- [47] Passreiter, M. et al. (1998) *J. Cell Biol.* 141, 373–383.
- [48] Schrader, M. et al. (1998) *J. Biol. Chem.* 273, 29607–29614.
- [49] Kalish, J.E. et al. (1996) *EMBO J.* 15, 3275–3285.
- [50] Chang, C.C. et al. (1997) *Nat. Genet.* 15, 385–388.
- [51] Okumoto, K. et al. (1998) *Mol. Cell. Biol.* 18, 4324–4336.
- [52] Elgersma, Y. et al. (1996) *J. Cell Biol.* 135, 97–109.
- [53] Erdmann, R. and Blobel, G. (1996) *J. Cell Biol.* 135, 111–121.
- [54] Gould, S.J. et al. (1996) *J. Cell Biol.* 135, 85–95.
- [55] Albertini, M. et al. (1997) *Cell* 89, 83–92.
- [56] Brocard, C. et al. (1997) *EMBO J.* 16, 5491–5500.
- [57] Komori, M. et al. (1997) *EMBO J.* 16, 44–53.
- [58] Fransen, M. et al. (1998) *Proc. Natl. Acad. Sci. USA* 95, 8087–8092.
- [59] Komori, M. et al. (1999) *FEBS Lett.* 457, 397–399.
- [60] Elgersma, Y. et al. (1997) *EMBO J.* 16, 7326–7341.
- [61] Eitzen, G.A. et al. (1997) *J. Cell Biol.* 137, 1265–1278.
- [62] Honsho, M. et al. (1998) *Am. J. Hum. Genet.* 63, 1622–1630.
- [63] South, A.T. and Gould, S.J. (1999) *J. Cell Biol.* 144, 255–266.
- [64] Huhse, B. et al. (1998) *J. Cell Biol.* 140, 49–60.
- [65] James, G.L. et al. (1994) *J. Biol. Chem.* 269, 14182–14190.
- [66] Kammerer, S. et al. (1997) *Genomics* 45, 200–210.
- [67] Götte, K. et al. (1998) *Mol. Cell. Biol.* 18, 616–628.
- [68] Matsuzono, Y. et al. (1999) *Proc. Natl. Acad. Sci. USA* 96, 2116–2121.
- [69] Snyder, W.B. et al. (1999) *Mol. Cell. Biol.* 10, 1745–1761.
- [70] Koller, A. et al. (1999) *J. Cell Biol.* 146, 99–112.